

Supplemental Table 1. Baseline characteristics of diabetes and non-diabetes

variable	Diabetes (n=549)	Non-diabetes(n=1478)	p-value
Age (years)	68.5±12.9	67.8±14.2	0.323
Male, n (%)	337 (61.4)	961 (65.0)	0.130
Smoke, n (%)	71 (12.9)	165 (11.2)	0.270
Hypertension, n (%)	291 (53.0)	682 (46.1)	0.006
Heart failure, n (%)	234 (42.6)	550 (37.2)	0.026
COPD, n (%)	2 (0.4)	18 (1.2)	0.084
Hyperlipidemia, n (%)	158 (28.8)	308 (20.8)	<0.001
Ventricular arrhythmia, n (%)	77 (14.0)	284 (19.2)	0.007
Cerebral infarction, n (%)	16 (2.9)	30 (2.0)	0.235
Admission drugs			
Aspirin, n (%)	461 (84.0)	1202 (81.3)	0.168
Clopidogrel, n (%)	345 (62.8)	900 (60.9)	0.423
Statin, n (%)	387 (70.5)	1020 (69.0)	0.521
β-blocker, n (%)	421 (76.7)	1167 (79.0)	0.270
ACEI, n (%)	324 (59.0)	890 (60.2)	0.624
Diuretic, n (%)	350 (63.8)	742 (50.2)	<0.001
Insulin, n (%)	184 (33.5)	190 (12.9)	<0.001
Oral hypoglycemic drugs, n (%)	74 (1.3)	11 (0.7)	<0.001
Laboratory data			
WBC, 10 <sup>9</sup> /L	11.6 (9.1-14.9)	11.6 (9.2-14.6)	0.968
Platelet count, 10 <sup>9</sup> /L	219 (166-271)	219.0 (177-272)	0.208
Hemoglobin, mg/dL	11.3 (9.8-12.5)	11.8 (10.4-13.4)	<0.001
Creatinine, mg/dL	1.1 (0.9-1.7)	0.9 (0.8-1.2)	<0.001
Urea nitrogen, mg/dL	23.0 (16.0-39.0)	17.0 (13.0-24.0)	<0.001
SOFA score	4.0 (2.0-7.0)	3.0 (1.0-5.0)	<0.001
Glucose_0, mg/dL	178.0 (140.0-251.0)	132.0 (111.0-160.0)	<0.001
Survival, n (%)	460 (83.8%)	1256 (85.0%)	0.508

COPD: chronic obstructive pulmonary disease; ACEI: angiotensin-converting enzyme inhibitor; WBC: white blood cell count; SOFA: sequential organ failure assessment.